What is claimed is:

1. Mixture of fibre reactive dyes comprising one or more dyestuffs of the formula (I)

and one or more dyestuffs of the general formula (II)

10

where

X¹, X² are independently a labile atom or group;

Ar¹ is an aromatic residue substituted by at least one -SO₃M group;

M is hydrogen or alkali metal, especially sodium;

15 Ar² is an aromatic radical substituted with at least one -SO₃M group;

a is 1 or 2 wherein,

if a is 2

L is a divalent radical typically of the form (y)

20

$$R^{1}$$
 $\stackrel{*}{\stackrel{N}{\nearrow}} L^{1}$ $\stackrel{*}{\stackrel{N}{\nearrow}} R^{2}$ (y)

where

 R^1 and R^2 are independently hydrogen, $C_1\text{-}C_4$ alkyl optionally substituted by $-\mathsf{OR}$, $-\mathsf{SR}$, $-\mathsf{SO}_3\mathsf{M}$ or X, or a phenyl group optionally substituted by a sulfonic acid group, - OR , $-\mathsf{C}_1\text{-}\mathsf{C}_4\text{-}$ alkyl, or NR'COR and $L^1 \quad \text{is arylene or alkylene optionally substituted by a sulfonic acid group, -OR, -C_1\text{-}C_4\text{-alkyl}$ –COOR, or NR'COR or

5

10


15

L is aminoethylpiperazine, under the proviso that if L is aminopiperazine, Ar¹ and Ar² are different or

if a is 1

L is a monovalent radical -NR³R⁴, -SR³ or -OR³ where

 R^3 and R^4 have one of the meanings of R^1 and R^2 or for -- NR^3R^4 , R^3 and R^4 can form a cyclic structure of the form (o)



20

25

where

U is an C_4 - C_6 alkyl residue optionally substituted by a substituent of formula Z and optionally interrupted by heteroatoms or heteroatom-containing groups such as -O-, $-NR^1$,

n is 1, 2 or 3 and

Z is hydrogen, optionally substituted $C_1\text{-}C_4$ alkyl, -OR 5 , -CO $_2\text{R}^5$, -COR 5 and

R⁵ is hydrogen, optionally substituted C₁-C₄ alkyl, optionally substituted vinyl, optionally substituted phenyl.

2. Dyestuff mixture according to claim 1

wherein

5

10

 X^1 and X^2 is independently chlorine, fluorine or 3 or 4-carboxypyridinium; Ar^1 and Ar^2 is independently a naphthyl residue substituted by at least one sulfo group (q-1)

where n is 1 to 3

or is a phenyl residue substituted by at least one sulfo group (q-2)

15 wherein

m is 1 or 2

p is 1 or 2 and

Y is independently hydrogen, halogen, R⁵, OR⁵, SR⁵, NHCOR⁵, where R⁵ is as given in claim 1.

20

25

3. Dyestuff mixture according to claim 1

wherein

X¹ and X² is chlorine:

Ar¹ and Ar² are independently a naphthyl residue of the formula (q-11) or (q-12)

$$MO_3$$
S

or

 SO_3M
 SO_3M
 $(q-11)$
 $(q-12)$

or are a phenyl residue substituted by at least one sulfo group (q-2) $(MO_3S)m$

wherein

m is 1 or 2

p is 1 or 2 and

Y is methyl.

10

15

20

5

- 4. Dyestuff mixture according to claim 1 wherein a is 1 and L is morpholine.
- 5.A dye mixture according to at least one of the claims 1 to 4 wherein a dye of formula (I) is present in the mixture in an amount of from 1% by weight to 99% by weight and a dye of the formula (II) is present in the mixture in an amount of from 99% by weight to 1% by weight.
- 6. A dye mixture according to at least one of the claims 1 to 4 wherein a dye of formula (I) is present in the mixture in an amount of from 10% by weight to 90% by weight and a dye of the formula (II) is present in the mixture in an amount of from 90% by weight to 10% by weight.
- 7. A process for preparing a dye mixture as claimed in one or more of claims 1 to 5, which comprises reacting chromophores of formula (III)

5

10

wherein Ar¹, Ar², X¹, X² and M are as defined in claim 1 with an appropriate mixture of 2-aminoethylpiperazine and a diamine H-L-H, or amine H-L, wherein L is as defined above, followed by precipitation using methylated spirits and conventional filtration.

8. A process for dyeing hydroxy- and/or carboxamido -cont aining fiber material, in which dyestuffs or dyestuff mixtures are applied to the material and the dyes are fixed to the material by means of heat or with the aid of an alkali or by means of heat and with the aid of an alkali, which comprises dye mixtures or dyestuffs as claimed in one or more of the claims 1 to 5.